

Cluster-state quantum computing enhanced by high-fidelity generalized measurements

D.N. Biggerstaff,¹ T. Rudolph,² R. Kaltenbaek,¹ D. Hamel,¹ G. Weihs,^{3,1} and K.J. Resch¹

¹*Institute for Quantum Computing and Department of Physics & Astronomy,
University of Waterloo, Waterloo, Canada, N2L 3G1*

²*QOLS, Blackett Laboratory, Imperial College London,
Prince Consort Road, London SW7 2BW, United Kingdom*

³*Institut für Experimentalphysik, Universität Innsbruck, Technikerstr. 25, 6020 Innsbruck, Austria*

We introduce and implement a technique to extend the quantum computational power of cluster states by replacing some projective measurements with generalized quantum measurements (POVMs). As an experimental demonstration we fully realize an arbitrary three-qubit cluster computation by implementing a tunable linear-optical POVM, as well as fast active feedforward, on a two-qubit photonic cluster state. Over 206 different computations, the average output fidelity is 0.9832 ± 0.0002 ; furthermore the error contribution from our POVM device and feedforward is only of $O(10^{-3})$, less than some recent thresholds for fault-tolerant cluster computing.

Measurement-based (cluster) computation [1] is an attractive alternative to standard circuit-based quantum computing. Instead of requiring multi-qubit gates, which are hard to implement experimentally, cluster computing requires only simple, single-qubit projective measurements. However, the prerequisite is a highly-entangled, multi-qubit cluster state. Thus far, laboratory cluster states [2, 3, 4, 5, 6, 7] have proven difficult to generate and limited in size. In order to make the most of these resources it is thus desirable to find means to extend the computational power of available clusters. Here we introduce one such technique, based on performing Positive Operator-Valued Measures (POVMs)[8, 9] on cluster-state qubits instead of standard projective measurements. As an experimental demonstration, we implement this technique to perform a three-qubit cluster computation for state preparation using linear optics and two entangled photons. Our results show the error introduced by the POVM apparatus and subsequent feedforward to be of $O(10^{-3})$, suggesting operation within recent thresholds for fault-tolerant cluster quantum computing [10].

Every POVM can be implemented by interacting an ancilla with the system to be measured, and performing projective measurement(s) on the combined Hilbert space [8]. Consider using the controlled-Z (CZ) operation (the interaction) to attach a new small cluster (the ancilla) to a qubit S (the system) in a cluster to be used for computation. By then performing single-qubit projective measurements on the ancillary cluster and S , we implement a POVM on S . For cluster computation, it matters only that the correct POVM is implemented; the manner in which this is accomplished is immaterial. In particular it may not be necessary to perform the (often technically challenging) task of interacting S with ancillary physical systems; the same POVM can be implemented using additional degrees of freedom of the qubit itself [11, 12], which are readily available in many architectures. For the photonic polarization qubits considered herein, one can employ an additional spatial-mode degree of freedom to implement an arbitrary 4-outcome POVM,




FIG. 1: (Color online): Cluster computing with POVMs. (a) A four-qubit ‘box’ cluster, one of the largest clusters achievable experimentally with optical qubits [2, 3, 4, 5, 6, 7]. By measuring the input cluster qubits and implementing the correct Pauli error correction, one achieves output equivalent to the circuit shown. Note: $\{X, Y, Z\}$: standard Pauli operators; H : Hadamard operator $(X+Z)/\sqrt{2}$; $Z_{(\beta)}$: $\exp(iZ\beta/2)$; α (on a cluster qubit): measurement in $|\alpha_{\pm}\rangle = |0\rangle \pm \exp(i\alpha)|1\rangle$. (b) A larger computation, and equivalent circuit, achievable via implementing POVMs on each input qubit of the same ‘box’ cluster. Cluster qubits with dashed outlines represent ‘virtual’ qubits simulated by performing POVMs $\{E\}_{\alpha}$ and $\{E\}_{\beta}$. (c) The computation we perform experimentally. By performing the POVM $\{E_m\}$ on the input qubit to our two-qubit cluster, we realize a three-qubit linear cluster computation, which in turn corresponds to a circuit for arbitrary pure state preparation via successive rotations of $|+\rangle$ around the z and x axes.

and all single-qubit POVMs can be obtained by suitable (classical) processing of such a POVM [11].

As a specific example, consider the 4-qubit box cluster [2], and equivalent circuit, shown in Fig. 1a). By performing POVMs on each of the first two qubits, one can effectively add ‘virtual’ qubits, thereby simulating a larger circuit, as shown in Fig. 1b).

Our approach differs fundamentally from recent experiments employing hyperentangled photon pairs for cluster computing [6, 7]. In contrast to those works, our method can replace arbitrarily large pieces of cluster, and avoids

the complications of sources producing particles entangled in multiple degrees of freedom. Furthermore our technique incorporates perfect automatic feedforward in the ‘virtual’ qubits, and is sufficiently versatile to serve as a useful primitive for large-scale cluster computers: POVMs (including those performed herein) can be applied, without modification, towards enhancing the computational power of *any* given cluster.

Before addressing the specifics of implementing an optical POVM to simplify cluster computations, let us mention some other closely related quantum information processing tasks. The ability to perform an arbitrary POVM on one qubit from a (not necessarily maximally) entangled pair constitutes the basic primitive of quantum steering [13], which underlies the optimal cheating attacks [14] on generalizations of the BB84 two-party bit commitment protocol. It also underlies the procedure for achieving maximum disturbance-free control [15], a basic primitive of quantum cryptography.

As a demonstration of our technique, consider the computation in Fig. 1c). We employ the smallest non-trivial cluster of two qubits; as depicted, a POVM on the first qubit allows a three-qubit computation. We label the ancillary qubit A and the system qubits S_1 and S_2 . In the manner of a standard cluster computation, we imagine A is initially in the state $|+\rangle_A$, is bonded to S_1 via a CZ gate CZ_{S_1A} , and is then projected into the basis $|\phi_{\pm}\rangle = |0\rangle \pm e^{i\phi}|1\rangle$. Depending on the sign of the outcome (+ or -), qubit S_1 must subsequently be projected into either $|\gamma_{\pm}\rangle$ or $X|\gamma_{\pm}\rangle$, respectively. These measurements are equivalent to the 4 projectors $\Pi_{ab}^{S_1A} = Z^a|\phi_{+}\rangle_A\langle\phi_{+}|Z^a \otimes X^a Z^b|\gamma_{+}\rangle_{S_1}\langle\gamma_{+}|Z^b X^a$, where $a, b \in \{0, 1\}$. We can represent this process as a POVM on S_1 as:

$$E_{ab}^{S_1} = \langle +_A | CZ_{S_1A} \Pi_{ab}^{S_1A} CZ_{S_1A} | +_A \rangle. \quad (1)$$

As can be readily verified, the POVM elements are $\{E_m\} = \frac{1}{2} \{\sigma_m |\chi\rangle\langle\chi| \sigma_m^\dagger\}$, where we use the index $m = 1, \dots, 4$, $|\chi(\phi, \gamma)\rangle = \cos(\phi/2)|0\rangle - ie^{i\gamma}\sin(\phi/2)|1\rangle$, and $\sigma_m \in \{\mathbb{1}, X, XZ, Z\}$.

The cluster model requires active feedforward in order to drive deterministic quantum computations despite inherently random measurement outcomes [4, 16]. When using POVMs on cluster qubits, the required feedforward depends on the measurement outcome: after performing POVM $\{E_m\}$ on S_1 and obtaining outcome m , the operator σ_m must be applied to the output qubit S_2 in order to recover the outcome of the circuit in Fig. 1c).

In our experiment we begin with photons in the Bell state $|\Phi^+\rangle_{S_1S_2}$, where $|\Phi^\pm\rangle = |HH\rangle \pm |VV\rangle$, and H and V indicate horizontal and vertical polarization, our $|0\rangle$ and $|1\rangle$, respectively. $|\Phi^+\rangle$ differs from a 2-qubit cluster by a Hadamard on one photon. We make a convenient adjustment of our second measurement angle so as to




FIG. 2: (Color online). Experimental implementation of cluster-state computing with a POVM. a) A source [17] produces maximally-entangled photon pairs coupled into single-mode fiber (SMF). b) The POVM is based on an optical interferometer constructed using calcite beam-displacers (BDs), which couple the polarization of the photon to the path, thereby enlarging the state space for the generalized measurement. Details are given in the text. c) Schematic of the cluster-state quantum computer. Photon-1 is measured in the POVM. Two Pockels cells (PCs), each fired dependent on the measurement outcome, actively perform the required correction of Pauli errors. A 50 m SMF serves to delay Photon-2, allowing time to trigger the PCs. The computational output is analyzed using quantum state tomography. Note: DM: dichroic mirror; IF: blocking and interference filter; PBS: polarizing beamsplitter; H(Q)WP: half (quarter) wave-plate; SPCM: fiber-coupled single-photon counting module.

implement the POVM:

$$\{\mathcal{E}_n(\phi, \theta)\} = \frac{1}{2} \{Z\rho^*Z, \rho^*, X\rho^*X, XZ\rho^*ZX\}, \quad (2)$$

where $\rho = |\psi\rangle\langle\psi|$, and $|\psi(\phi, \theta)\rangle = \cos(\phi/2)|H\rangle + e^{i\theta}\sin(\phi/2)|V\rangle$. After performing $\{\mathcal{E}_n\}$ on photon S_1 from our actual 2-photon state $|\Phi^+\rangle_{S_1S_2}$, obtaining outcome n , and implementing $\sigma_n \in \{Z, \mathbb{1}, X, XZ\}$ on S_2 , the output will be state $|\psi(\phi, \theta)\rangle$, where ϕ and θ are adjustable experimental parameters which respectively correspond to the polar and azimuthal angle of the output in the Bloch sphere representation.

We generate entangled photons as shown in Fig. 2a) [17]. A grating-stabilized diode laser outputs 0.86 mW at 404.5 nm to bi-directionally pump a 25 mm periodically-poled KTiOPO₄ (PPKTP) crystal in a polarization-dependent Sagnac interferometer, yielding 809 nm entangled photons via type-II parametric downconversion. At this power, the SMFs typically output singles (coincidence) rates of 150 (30) kHz. Polarization controllers (bat-ears) in the SMFs ensure the output is $|\Phi^+\rangle$.

The apparatus for performing the POVM $\{\mathcal{E}_n\}$ is depicted schematically in Fig. 2b). It is a polarization-

based double interferometer employing calcite beam displacers (BDs). Due to transverse walk-off, these couple polarization with optical path, enlarging the state space from dimension two to four. The settings $\{\phi, \theta\}$ of the POVM are determined by half-wave plates (HWPs) in the interferometer. Polarization measurements are implemented using polarizing beamsplitters (PBSs). The four output modes are coupled into SMF and detected using single-photon counting modules (SPCs). The BD construction is inherently phase-stable because the interfering paths propagate through common optics [18]. Furthermore, we align the setup using an 809 nm diode laser injected through the input SMF and a removable polarizing optic, and typically measure classical interference visibilities $> 99.8\%$; thus the setup is promising for high-fidelity, stable operation.

The action of this apparatus can be understood as follows. An arbitrary input qubit is in state $a|H\rangle + b|V\rangle$. The first BD displaces the H - relative to the V -component, introducing a ‘path qubit’ with basis states upper $|U\rangle$ and lower $|L\rangle$, and thereby creating the entangled state $a|HU\rangle + b|VL\rangle$. The polarization qubit is then measured using a HWP at angle $\phi/4$ and a PBS. HWPs at 45° then flip the polarization in path U (L) in the transmitted (reflected) arm. HWPs at 0° are included in the other path to balance path-lengths, and allow the phase to be adjusted via tilting about their vertical axes. Recombining the paths at subsequent BDs converts the ‘path’ qubit back to polarization; this qubit is then measured via a QWP at 45° , a HWP at $-\theta/4$ ($+\theta/4$), and a PBS in the transmitted (reflected) arm. This yields the POVM elements $\{\mathcal{E}_n\}$ as follows: The outcome TR which stems from (T)ransmission at the first PBS and (R)eflection at the second then corresponds to $\rho^*(\phi, \theta)/2$, and the TT, RT and RR outcomes to $Z\rho^*Z/2$, $X\rho^*X/2$, and $XZ\rho^*ZX/2$, respectively.

We implement the necessary feedforward using two fast RbTiOPO₄ (RTP) Pockels cells (PCs) (Leysop RTP4-20-AR800), able to switch to their half-wave voltage of 1.027 kV in < 5 ns, both oriented such as to enact an X operation (i.e. a HWP at 45°) when triggered. The first PC will be triggered by POVM outcomes RT or RR; the second is surrounded by HWPs which rotate its action to Z , and is triggered by outcomes TT or RR. Photon-2 is stored in a 50m SMF for 250 ns of delay to allow ample time for detection, logic, and triggering the PCs. Note that one stage of feedforward is incorporated directly into the design of the POVM since the angle of the last HWP is dependent on the outcome of the polarization measurement. This reduces experimental cost and complexity, as we require one less PC over a direct implementation [4]. It also improves computational speed, as each additional PC requires delaying the relevant photon by $O(100)$ ns to allow time for detection, logic, and triggering; in our POVM this feedforward requires a mere couple ns of optical path. After correction, we perform state tomography




FIG. 3: (Color online:) Experimentally reconstructed density matrix $\rho_{S_1S_2}$ of our 2-photon cluster state: real part (left) and imaginary part (right). The fidelity with the ideal state $|\Phi^-\rangle$ is $F = 0.980 \pm 0.001$; the tangle is $T = 0.926 \pm 0.002$ and the purity, $\text{Tr}(\rho^2)$, is $P = 0.963 \pm 0.002$ [19, 20].

on Photon-2 using a HWP, QWP, PBS, and SPCM. Our raw data consists of coincidence counts between this output and any one of the four POVM outcomes.

We characterized the entangled state generated in fibers 1 and 2 (see Fig 2a) via over-complete state tomography. To analyze Photon-1, we employed the first two waveplates in the POVM and the H output of the first BD; the V output was blocked, and the other waveplates set so as to direct all photons to output TT. We then counted coincidences between this output and that of the polarization analyzer following the (switched off) PCs. The density matrix is reconstructed via a maximum-likelihood technique [21], and shown in Fig. 3. Our measured state $\rho_{S_1S_2}$ has fidelity $F = 0.980$ with $|\Phi^-\rangle$. When not in use for tomography, the QWP at the beginning of the POVM is removed and the HWP set to 0° , which maps the $|\Phi^-\rangle$ source state to $|\Phi^+\rangle$.

We tested this cluster computer by performing computations with 206 different measurement settings $\{\theta, \phi\}$ over a period of 4 hours. The target output states $\{|\psi(\phi, \theta)\rangle\}$ include the six eigenstates of X , Y and Z , and 200 numerically-generated settings designed to be spread evenly over the surface of the Bloch sphere. For each computation, the output density matrix ρ_m is tomographically reconstructed based on coincidence measurements integrated over 8 s for each of six analyzer settings (the eigenstates of X , Y and Z .) We record maximum coincidence rates of about 3 kHz, summed over the four outputs.

Using the reconstructed density matrix $\rho_{S_1S_2}$ (Fig. 3), and assuming a perfect POVM and feedforward, we can estimate the output of the computation: $\rho_e(\phi, \theta) = \sum_{n=1}^4 \sigma_n \text{Tr}_1[(\mathcal{E}_n \otimes \mathbb{1})\rho_{S_1S_2}(\mathcal{E}_n \otimes \mathbb{1})^\dagger] \sigma_n^\dagger$. Fig. 4a) shows the fidelity between ρ_e and the target output $|\psi\rangle$.

Fig. 4b) shows the fidelity $F(\rho_m, |\psi\rangle)$ between our measured states and the target outputs. For these 206 computations the mean fidelity is 0.9832 ± 0.0002 , where the uncertainty is the standard error in the mean. This compares favorably with the expected mean fidelity




FIG. 4: (Color) Expected and measured fidelity of computational output. a) The expected output fidelity based on the measured cluster state, $\rho_{S_1 S_2}$, assuming perfect POVM operation and feedforward. b) The measured output fidelity with the target outputs; the mean is $F = 0.9832 \pm 0.0002$. c) The deviation from unity of the fidelity between the expected and measured states $1 - F(\rho_e, \rho_m)$; the mean is $(1 - F) = (1.16 \pm 0.05) \times 10^{-3}$. This shows our POVM and feedforward to be operating with very high fidelity.

$F(\rho_e, |\psi\rangle) = 0.9865 \pm 0.0001$ for the same 206 states. The distribution of $F(\rho_m, |\psi\rangle)$ is largely explained by two factors: expected variance due to the imperfect entangled state, and fluctuations from Poissonian counting statistics, as determined by a Monte Carlo simulation.

We characterize the errors introduced by the POVM and feedforward using the quantity $(1 - F(\rho_e, \rho_m))$, shown in Fig. 4c), where F is the mixed-state fidelity [20]. With a mean error of only $(1 - F) = (1.16 \pm 0.05) \times 10^{-3}$, our results demonstrate remarkable agreement with the model (Fig. 4a) based on $\rho_{S_1 S_2}$. More importantly, this shows that our POVM and feedforward are very stable and exhibit a low error rate, comparable to some thresholds for fault-tolerant cluster computing [10].

We have shown how POVMs may be employed in cluster quantum computing to increase the computational power of any given cluster. Furthermore we have experimentally demonstrated this technique by fully realizing

an arbitrary, high-fidelity three-qubit cluster computation using two photons. Some feedforward steps can be incorporated into the design of the POVM, significantly improving computational speed while reducing experimental complexity. This technique should be incorporated into future cluster computers to maximize the utility of available resources.

We thank N. Killoran, N. Lütkenhaus, and K.M. Schreier for valuable discussions, and Z.-W. Wang for designing and building our TTL logic. D.B. acknowledges financial support from the Mike and Ophelia Lazaridis Fellowship. T.R. acknowledges support from EPSRC and the US Army Research Office. We are grateful for financial support from NSERC, OCE, and CFI.

-
- [1] H.J. Briegel and R. Raussendorf, Phys. Rev. Lett. **86**, 910 (2001); R. Raussendorf and H.J. Briegel, *ibid.* **86**, 5188 (2001); R. Raussendorf, D.E. Browne, and H.J. Briegel, Phys. Rev. A **68**, 022312 (2003).
 - [2] P. Walther *et al.*, Nature **434**, 169 (2005).
 - [3] N. Kiesel *et al.*, Phys. Rev. Lett. **95**, 210502 (2005).
 - [4] R. Prevedel *et al.*, Nature **445**, 65 (2007).
 - [5] C.-Y. Lu *et al.*, Nature Physics **3**, 91 (2007).
 - [6] K. Chen *et al.*, Phys. Rev. Lett. **99**, 120503 (2007).
 - [7] G. Vallone *et al.*, Phys. Rev. Lett. **98**, 180502 (2007); G. Vallone *et al.*, Phys. Rev. A **78**, 042335 (2008).
 - [8] A. Peres, *Quantum theory: concepts and methods* (Dordrecht: Kluwer, 1993), ISBN 0792325494.
 - [9] S.M. Barnett and E. Riis, J. Mod. Opt. **44**, 1061 (1997); S.E. Ahnert and M.C. Payne, Phys. Rev. A **71**, 012330 (2005).
 - [10] R. Raussendorf, J. Harrington and K. Goyal, New J. Phys. **9**, 199 (2007).
 - [11] J. Preskill. "Lecture notes on quantum information and computation", www.theory.caltech.edu/people/preskill/ph229/#lecture.
 - [12] B.P. Lanyon *et al.*, Nat. Phys. **5**, 134 (2009); T.C. Ralph, K.J. Resch and A. Gilchrist, Phys. Rev. A **75** 022313 (2007).
 - [13] E. Schrödinger, Proc. Camb. Phil. Soc. **31**, 555 (1935); L.P. Hughston, R. Jozsa and W.K. Wootters, Phys. Lett. A **183**, 14 (1993).
 - [14] C.H. Bennett and G. Brassard, Proc. IEEE International Conf. on Computers, Systems, and Signal Processing, IEEE (New York), 175 (1984); R.W. Spekkens and T. Rudolph, Quantum Inform. Compu. **2**, 66 (2002).
 - [15] T. Rudolph and R.W. Spekkens, Phys. Rev. A **70**, 052306 (2004).
 - [16] T.B. Pittman, B.C. Jacobs, and J.D. Franson, Phys. Rev. A **66**, 052305 (2002).
 - [17] T. Kim, M. Fiorentino, and F.N.C. Wong, Phys. Rev. A **73**, 012316 (2006); A. Fedrizzi *et al.*, Optics Express **15**, 15377 (2007).
 - [18] J.L. O'Brien *et al.*, Nature **426**, 264 (2003).
 - [19] A.G. White *et al.*, Phys. Rev. A **65**, 012301 (2001).
 - [20] Definition: $F(\rho, \sigma) = [\text{Tr}(\sqrt{\sqrt{\rho}\sigma\sqrt{\rho}})]^2$. R. Jozsa, J. Modern Optics **41**, 2315 (1994).
 - [21] D.F.V. James *et al.*, Phys. Rev. A **64**, 052312 (2001).